

Hair Transplant Instructions – Before Your Hair Transplant

Time	Hair Care		Instructions
6 weeks before	Grow your hair to be at least 1 inch long by your hair transplant day and preferably 2 inches long	Do scalp stretching exercises if instructed by Dr. Simmons	Do lab tests ASAP
3 weeks before	Do not cut your hair shorter than 1 inch long You can start Minoxidil to jump start new hair growth and reduce postoperative shedding		Must do lab tests Limit green leafy vegetables to 1 salad per day and don't eat lots of nuts Stop multivitamins and Vitamin E and Herbal supplements/meds
2 weeks before	Keep your hair at least 1 inch long		Do lab tests Pay balance
1 week before	Stop using Minoxidil but you can continue taking Finasteride if you take it normally You can dye your hair right up to the day before your hair transplant	No vigorous exercise, cardio, or weights	No Alcohol No Aspirin, or Anti-inflammatories like Advil or Motrin but Tylenol is OK Arrange for someone to pick you up after your hair transplant Get a travel pillow, firm hat that does not touch the top of your head (like a trucker's hat), and buy 2% minoxidil lotion, if you are going to start it after your hair transplant
1 day before	Wash your hair with shampoo at night then again in the morning of your hair transplant		Pack any music ipod, cds, mp3 player, or DVDs that you want (but we have music and DVDs) Pack a full-button or full-zipper shirt that does not have to be taken off over your head Confirm your transportation home Check the weather and set an alarm so that you can be sure to arrive by 7:30 am
Before you leave for your hair transplant	Wash your hair wi th shampoo but don't use conditioner or hair products		Eat breakfast but only decaf coffee Bring your hat, full-button or full-zipper shirt, and music/DVDs If someone is going to pick you up, make sure they will be available any time between 4 pm and 10 pm and ask them to call us for an update at 416.287.3733 at 2 pm You cannot answer your blackberry or phone today



Hair Transplant Instructions – After Your Hair Transplant

Your Hair Transplant Day

We will keep you safe. You will feel minor (2-3/10) discomfort when we make the areas numb but should expect a calm and comfortable day. You will spend most of your day watching movies, TV, or listening to music but will have a lunch and bathroom breaks. Save any postoperative questions for the end of the day. We will review all of the instructions and answer questions and give you medications before you leave. We can't predict exactly how long your hair transplant will take so your travel arrangements must be flexible.

	What to Expect	Hair Care	Activities	Instructions
The first night (Day 0)	The recipient area will not hurt You may have swelling in your forehead and in the corners of your eyes from the local anaethetics You may feel tightness, pressure, throbbing, or aching around the sutures. Tonight will be the most difficult night so take pain medications early to prevent severe pain before it develops	Do not touch the grafted area or get it wet Leave your dressing on overnight No Minoxidil Continue Finasteride (Propecia/Proscar) throughout your recovery period, if you take it normally	No driving tonight No alcohol tonight Rest and sleep with your head Elevated 30 degrees. Putting pillows against the firm arm-rest of a couch or sleeping in a recliner chair is better than using extra pillows in bed. A travel pillow reduces pressure on the sutures for more comfort and keeps you from rolling in bed a lot Apply Ice to your forehead 10 minutes out of every waking hour To prevent swelling remember RIPE = Rest, Ice, Prednisone, and Elevation	For 1/10 to 3/10 pain take 1 Percocet with food. Take 1 more 1 hour later if not relieved. For 4/10 to 6/10 pain take 2 Percocet tablets with food. If you have 7/10 pain or higher or you have taken 2 Percocets and didn't get relief, phone Dr. Simmons at 416.287.3733 Do not text or email. Do Phone. Take Gravol every 6 hours as needed for nausea. Take Restoril before bed to help sleep.
Days 1 and 2	Your forehead and the corners of your eyes may have swelling or bruising You may still need pain tablets in the morning for the donor area but the tightness usually eases off as the day goes on.	Do not touch the grafted area or get it wet Take off the dressing in the morning of Day 1 Wash the donor area with a washcloth and shampoo Apply ointment once daily to the donor (sutured) area No Minoxidil	Rest and sleep with your head Elevated 30 degrees To prevent swelling remember RIPE = Rest, Ice, Prednisone, and Elevation	Take Prednisone with breakfast unless you have stomach pain or Dr. Simmons said otherwise Take Percocet, Gravol, and Restoril as needed You can resume herbals, supplements, and Vitamins



Hair Transplant Instructions – After Your Hair Transplant

	What to Expect	Hair Care	Activities	Instructions
Days 3 and 4	You may have noticeable swelling around your eyes that can even make your vision blurry temporarily	You can rinse the transplanted area in the bathtub by pouring water & shampoo then water alone from a cup onto an area in front or behind the transplanted area and letting it flow over the transplanted area. Do not rub the transplanted area or try to style your hair. Apply ointment once daily to the donor (sutured) area Let your hair dry naturally You can spray on a saltwater mixture frequently if your scalp feels dry or itchy. You would add ½ tsp salt or seasalt to 8 ounces of water or distilled water and use a gentle mist sprayer You can start using Minoxidil 2% (available without a prescription) twice daily but should not rub it into your scalp (see other handout)	You can sleep normally from Day 3 onward and should stop using sleeping pills Avoid cardio	Take your last Prednisone on Day 3 Take Tylenol, Advil as needed and only use Percocet if not relieved with those medicarions
Days 5 and 6	Day 5 will probably be the last day you have swelling around your eyes and by Day 6 it may have moved down into your cheeks A few scabs may come off and some short and long hairs may come with them but that does not mean you are losing grafts. Grafts are secure and will only come out with forceful rubbing, picking, or trauma	Wash your whole head in the bath once daily by pouring shampoo and water from a cup and rubbing in gentle circles then rinsing the sane way Apply ointment once daily to the donor (sutured) area Let your hair air-dry You can gently style your hair with your hands	You can resume cardio but shouldn't lift weights	



Hair Transplant Instructions – After Your Hair Transplant

Days 7, 8, and 9	Your swelling will be mostly	Wash your whole head	Cardio is OK but not weights or	
	gone	gently in the shower but	contact sports	
	Your recipient area will still	don't let the jets hit the		
	have scabs and may have little	recipient area directly.		
	bald rivers or cracks in the	Apply ointment once daily to		
	crust	the donor (sutured) area		
	You may shed hairs and scabs	Let your hair air-dry		
	but won't lose grafts	Gently style your hair with		
	_	your hands		

Day 7 to 13 (most often 10 or 11)

Suture Removal

	What to Expect	Hair Care	Activities
Days 10 to 13	Your skin may be pink at first. You can use	You can shower normally	You can do cardio exercise
	your hair or non-comedogenic makeup to	Rub the remaining crusts off firmly in the	You can swim and dive
	conceal it.	shower but don't pick them	No weights or contact sports
	Some stubble may remain after the scabs	Apply baby oil or ointment to the grafted	
	come off but some will shed	area when the skin is still wet and leave it in	
	You will shed short and long hairs but will	You can shower every 3 hours as needed	
	not lose grafts	until the crusts are all off	
		You can use all hair products	
		You can use a hair dryer	
		Continue Finasteride/Propecia/Proscar if	
		you take it normally	
		Use Minoxidil twice daily if you have	
		chosen to use it. Stop it if your skin is very	
		pink or itchy or you get pimples	
Days 14 to 27	You can experience post-operative shedding	Usual hair care	You can do almost all activities including
	Pinkness will usually be disappearing but can	Don't cut or dye your hair yet	weight lifting, contact sports, and wearing
	persist for some	Continue Finasteride/Propecia/ Proscar and	helmets
		Minoxidil unless minoxidil causes pinkness or	Do not stretch or strain your scalp or neck
		pimples	until 3 months after your hair transplant
			Call 416.287.3733 for an appointment if you
1 . 2 M . 4	D' 1 (T) 1 (1		get a lot of pimples
1 to 3 Months	Pimples may start. They may be tender or	You can cut and dye your hair	You can do almost all activities
	cause bleeding when you rub your scalp in	Hair stylist should look before cutting so your	Do not stretch or strain your scalp or neck
	the shower.	hides the donor	until 3 months after your hair transplant
	Your hair will typically look the same or	scar	Call 416.287.3733 for an appointment if you
	worse than you started	Continue Propecia/ Proscar and Minoxidil	get a lot of pimples



Hair Transplant Instructions - After Your Hair Transplant

You will likely shed almost all of the transplanted stubble in the first 2 – 6 weeks after your hair transplant and won't see new hair until about 4 months after your hair transplant. You will not see daily improvement but will see big monthly changes until about 8 months then subtle changes from then until 1 year after your hair transplant. It is best to leave your hair a bit longer than usual to get the most out of your new hair.

Your donor scar will be pink for 6 months and you may get some pimples in the donor area for the first 6-9 months. The hair is trimmed above and below the incision and you may get some shedding of hair around your incision. You should always ask your hair stylist to look at the scar first and to avoid cutting your hair too short.

In the first few months you may experience pinkness, pimples, shock loss, numbness, or itchiness but won't experience all of them. There are ways to cope with each of these concerns and you should contact Dr. Simmons if you are having difficulties.

We will call you for follow-up appointments or emails at:

1 month to look for and remedy nuisance side effects 6 months to check for early but incomplete results 12 months to check your final results

Please call us at 416.287.3733 anytime with any concern. Please do not email or text if you have urgent concerns.